

1917

# Studies in the genus Hydnocera Newman: the species of the New England states /

Edward A. Chapin

*University of Massachusetts Amherst*

Follow this and additional works at: <https://scholarworks.umass.edu/theses>

---

Chapin, Edward A., "Studies in the genus Hydnocera Newman: the species of the New England states /" (1917). *Masters Theses 1911 - February 2014*. 1237.

Retrieved from <https://scholarworks.umass.edu/theses/1237>

This thesis is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Masters Theses 1911 - February 2014 by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

Studies in the Genus *Hydnocera* Newnan.  
The Species of the New England States.

By Edward A. Chapin, Ph. B.

Thesis  
submitted for the degree of  
Master of Science.

Massachusetts Agricultural College,  
Amherst, Massachusetts.

May 15, 1917.

595  
.97  
C367  
ENTOMOLOGY

Studies in the Genus *Hydnocera* Newm., (Col.).

The Species of the New England States.\*

Edward A. Chapin, Ph. B.

Introduction.

The following paper comprises the partial results of a systematic study of the insects of the genus *Hydnocera* Newman of the coleopterous family Cleridae and is intended as a preliminary to a complete systematic revision of the genus for the United States. The greater part of the work was done at the Massachusetts Agricultural College during the past year and is offered as fulfillment in part of the requirements for the degree of Master of Science. During the course of the work it has been the good fortune of the writer to receive for study the entire collection of Mr. Albert B. Wolcott of the Field Museum of Natural History, Chicago, Ill.; the general collections of the Massachusetts Agricultural College, including the major part of the Cleridae from the collection of the late Mr. L. H. Joutel; Kansas Agricultural College; Kansas State University; University of Maine; Montana Agricultural Experiment Station and other smaller lots. In addition to these, the author has seen and studied the material in the Leconte and Blanchard collections at Harvard University, the material in the collection of the Boston Society of Natural History, and the New England material at the American Museum of Natural History, New York City. Whenever possible, the type itself

has been studied and drawings included in the plate made from

\* Contribution from the Entomological Laboratory of the Massachusetts Agricultural College, Amherst, Massachusetts.





such a source have been noted.

At this time it gives me great pleasure to record my thanks to Drs. H. T. Fernald and G. C. Crampton, both of whom have assisted me in all possible ways. To Mr. Albert B. Wolcott, I offer my thanks and gratitude, not only for the material placed at my disposal but for the countless favors in the way of literature and suggestions which he has at all times offered most freely and generously. Thanks are also due Dr. W. S. Regan, Messrs. Nathan Banks of the Museum of Comparative Zoology, Harvard, C. W. Johnson of the Boston Society of Natural History, A. J. Mutchler of the American Museum of Natural History, New York and to all others who have so kindly placed material at my disposal.

#### Historical.

Subfamily Hydnocerini : The genus Hydnocera Newm. falls naturally into the fourth of the six subfamilies of the Cleridae and is itself the type genus of that subfamily, the Hydnocerini. This group of genera is separated from the rest of the family by the following characteristics, the arrangement of which is adapted from Mr. Sigmund Schenkling's work on the family in the Genera Insectorum.

- |  |                        |
|--|------------------------|
| Tarsi of five segments; thorax without lateral margins. . . . .                    | 1.                     |
| Tarsi of four segments; thorax with more or less distinct lateral margins. . . . . | 4.                     |
| 1. All tarsal segments visible from above. . . . .                                 | <u>Tillini</u> .       |
| First segment covered from above by second, sometimes minute. . . . .              | 2.                     |
| 2. Eyes entire or only minutely emarginate at base of antennae . . . . .           | <u>Hydnocerini</u> .   |
| Eyes distinctly emarginate. . . . .  | 3.                     |
| 3. Eyes emarginate in front, sometimes but slightly. . . . .                       | <u>Clerini</u> .       |
| Eyes emarginate on inside. . . . .   | <u>Phyllobaenini</u> . |
| 4. Antennae serrate or with three large terminal segments. . . . .                 | <u>Enopliini</u> .     |
| Antennae with a small triarticulate club. . . . .                                  | <u>Corynetini</u> .    |

The first of these is the fact that the United States is a young nation, and that its history is a history of growth and expansion. The second is the fact that the United States is a nation of immigrants, and that its history is a history of the struggle for a common identity. The third is the fact that the United States is a nation of free men, and that its history is a history of the struggle for freedom and justice.

CHAPTER I

The first of these is the fact that the United States is a young nation, and that its history is a history of growth and expansion. The second is the fact that the United States is a nation of immigrants, and that its history is a history of the struggle for a common identity. The third is the fact that the United States is a nation of free men, and that its history is a history of the struggle for freedom and justice.

CHAPTER II

The first of these is the fact that the United States is a young nation, and that its history is a history of growth and expansion. The second is the fact that the United States is a nation of immigrants, and that its history is a history of the struggle for a common identity. The third is the fact that the United States is a nation of free men, and that its history is a history of the struggle for freedom and justice.



In 1903, Schenkling assigned the following genera to the subfamily Hydnocerini : *Isolemidia* Gorham, *Hydnocera* Newman, *Ellipotoma* Spinola, with possibly *Theano* Castelnau. In 1910, C. J. Gahan, of the British Museum, basing his work on the great collections of that institution, made one change in the above arrangement, that of relegating *Ellipotoma* Spinola to the *Enopliini*. Still later, in 1912, A. B. Wolcott took the genus *Zenodosus* Wolcott from the *Clerini* where it had been placed since about 1830 and added it to the genera of the *Hydnocerini*. Of the genera now considered as belonging in this group, but two are to be found in the United States, one *Zenodosus*, *Z. sanguineus* Say, widely distributed over North America is the sole member of the genus. The other genus, *Hydnocera* Newm., forms the subject of this paper.

Genus *Hydnocera* Newman : The first insect to be described, which today is included within the confines of this genus, was the species - *humeralis* - described by Thomas Say in the year 1823 as *Clerus humeralis*. Three more species were described by the same author before 1838, when Newman erected a new genus *Hydnocera* to include the four species of Say, *humeralis*, *pallipennis*, *verticalis* and *unifasciata*, as well as three new ones of his own making, *rufipes*, *curtipennis* and *aegra*. At the time, Newman published the description of a species which he designated as *serrata* and by page priority, this is the type of the genus. But as *serrata* is a synonym of *pallipennis* Say, the latter becomes the type. Four years later, Klug, in his attempt at a synopsis of the *Cleridae* of the world, refused to recognise *Hydnocera* and placed under the head of *Clerus*, species that are





now considered as belonging to no less than fifteen genera! However, in 1844, Maximilian Spinola published a masterpiece on the same subject and reestablished Newman's genus, adding some new species and noting that the species are solely North American in distribution. Five years later two interesting papers appeared, one by Mr. Adam White entitled "A Catalogue of the Coleoptera in the British Museum, Part IV, Cleridae". Here the author disregards Spinola's note and places in Hydnocera, seven forms from Australia. The second paper concerns us more as it marks the first attempt to tabulate the Cleridae of the United States, made by Dr. John L. LeConte. Fourteen species of the genus Hydnocera are here noted, of which eleven are now considered valid, although one, an actual new species was wrongly identified and placed as a previously described form. Since that time many new species have been added to the list, the greater part of them appearing in the Biologia Centrali-Americana, under the authorship of the Rev. H. S. Gorham. Other workers who have added to the list are A. B. Wolcott, H. F. Wickham, H. C. Fall, Charles Schaeffer, F. C. Van Dyke and the author. The present list of recognised forms numbers 120, of which eleven are to be found in the New England States.

#### External Anatomy.

Form varies from very elongate with parallel sides to quite robust with the lateral margins of the elytra decidedly attenuate toward the posterior. Size quite variable. In H. bicolor Lec. and H. pusilla Schaeff., the length hardly exceeds two and one-half millimeters while in some specimens of H. longa Lec. the length may reach a maximum of seven millimeters. Vestiture. All





species are clothed with two types of pubescence to a greater or lesser degree. The pubescence of the first type is, so far as I am able to determine, universally present and consists of long, erect, sparsely distributed hairs, evenly scattered over the entire body, with the exception of the mouth-parts and tarsi. the second type, found to some extent on nearly all the species which can be considered as typical and especially in the case of such as scabra Lac. and subfasciata Lec., and consists of short, depressed, more or less scale-like hairs, usually pale or silvery in color.

Head. Exclusive of appendages, consists of a single sclerite, all cervical sutures being obliterated. Front large and flat, nearly equilateral but in some cases trapezoidal in shape. Lateral margins parallel or convergent toward labrum, sharply defined. At upper part passing imperceptibly into the vertex, which is short and broad. Both front and vertex carry, in all species, punctures. The coarseness and density of the punctures constitutes a character of use in the separation of species. Eyes usually large, quite convex, the inner margin straight, rarely convex. Surface is finely granulated with always a few long hairs arising from between the facets. The prominence of the eyes is always lateral, never in front of the face, and usually causes the head to appear wider than the elytra at the humeri. Antennae short, filiform and capitate, arising at the lower part of the anterior border of the eye (at which point the eye carries a minute emargination), composed of eleven segments. The first segment is large, cylindrical but somewhat arcuate, and is usually about the length of the terminal club. The second segment is considerably shorter than the first or third,

The first of these is the fact that the  
 system is not a simple one, but a  
 complex one, involving many factors  
 which are not easily understood or  
 explained. The second is the fact  
 that the system is not a simple one,  
 but a complex one, involving many  
 factors which are not easily under-  
 stood or explained. The third is the  
 fact that the system is not a simple  
 one, but a complex one, involving  
 many factors which are not easily  
 understood or explained. The fourth  
 is the fact that the system is not a  
 simple one, but a complex one, in-  
 volving many factors which are not  
 easily understood or explained. The  
 fifth is the fact that the system is  
 not a simple one, but a complex one,  
 involving many factors which are not  
 easily understood or explained. The  
 sixth is the fact that the system is  
 not a simple one, but a complex one,  
 involving many factors which are not  
 easily understood or explained. The  
 seventh is the fact that the system  
 is not a simple one, but a complex  
 one, involving many factors which are  
 not easily understood or explained. The  
 eighth is the fact that the system is  
 not a simple one, but a complex one,  
 involving many factors which are not  
 easily understood or explained. The  
 ninth is the fact that the system is  
 not a simple one, but a complex one,  
 involving many factors which are not  
 easily understood or explained. The  
 tenth is the fact that the system is  
 not a simple one, but a complex one,  
 involving many factors which are not  
 easily understood or explained.



and is more globose in form, in some species being nearly spherical. The third segment is cylindrical, very long and thin, usually as long as the first but much less thick. Segments four to nine are much the same in shape as the third, but each becoming shorter and broader than the preceeding. The tenth segment is very large and globose but in some species is somewhat flattened but always forming with the eleventh and last segment, a distinct club. The terminal segment is very small in most cases and the suture is usually so disguised by the fine pubescence as to be invisible. The color varies from testaceous to piceous but is never pigmented with metallic colors, as is the body in some cases. Pubescence usually fine and moderately dense. Labrum transverse, trapezoidal in form, separated from the front by a more or less distinct suture, anterior margin entire or very slightly emarginate, pubescence very sparse but evenly distributed. Mandibles slender falciform, decidedly acuminate toward tips, quite large, but when closed concealed beneath the labrum. On the inner side each bears one or more teeth near the middle, the larger tooth (when there are more than one) or the single tooth is placed slightly basal to the middle and is comparable to the retinaculum of the Carabidae. The mandibles never carry pubescence of either type. Maxillae moderately large, both lobes present. The outer lobe or lacinea is oblong oval in form, slightly broader at the distal end. The entire outer surface is covered with rather dense, moderately coarse hairs, while the margin is set with stout spiniform setae. The inner lobe or galea is more narrow but quite as long, with the lateral margins parallel and the tip rounded; the vestiture is similar to that of the lacinea but with spiniform setae only at the rounded end. Maxillary palpi com-





posed of four segments. Of these, the first and second segments are similar in form and size, short and stout, together nearly as long as the galea. Pubescence sparse, consisting of a few long hairs placed at irregular intervals. The third segment is cylindrical, somewhat longer than the second or first and in some species is slightly fusiform. Terminal segment about two-thirds as long as the preceeding, somewhat conical in form. The pubescence of the last two segments similar to that of the first. Labium small, quadrate, not separated from the mentum by a distinct suture, moderately densely clothed with long hairs. Labial palpi of three segments, arising from the lateral margins of the labium slightly behind the middle. Segments one and two long, thin and cylindrical, sparsely pubescent. Terminal segment very large, triangular and somewhat flattened. Pubescence moderately dense but short and indistinct. Ligula and paraglossae not distinct from the labium. Gula sutures short, parallel, enclosing the basal part of the mentum.

Thorax composed of three distinct regions. Prothorax of one piece, all sutures between the notum, pleurae and sternum being obliterated. In the species that are nearest like the type, the prothorax is distinctly broader than long, with a conspicuous lateral dilation slightly anterior to the middle. Anterior to this dilation there is a rather deep, transverse impression which is most conspicuous on the disk and which becomes nearly obsolete under the side prominences. Near the base of the thorax, there is a second transverse impression somewhat similar to the first, but usually broader and more conspicuous at the sides. In the group of species of which curtipennis is typical, the thorax is much longer than broad, nearly cylindrical with very





weak lateral dilation. The anterior transverse impression is much less evident while the basal impression is quite strong. The surface of the entire thorax of the first type is covered with rather densely placed, moderately fine to coarse punctures, while those of the second type have the thorax nearly smooth with a few scattered coarse punctures. Anterior coxal cavities wide open behind, in most species nearly confluent. Mesothorax very short and inconspicuous, composed of the normal number of sclerites. Notum concealed, partially membranous. Sternum small, transverse, truncate on the anterior margin, on the posterior margin prolonged into a short quadrate process which extends between the median coxae, thus causing them to be slightly separated. The pleurae are each divided by a longitudinal suture into two parts, the epimeron and episternum. Both pieces are somewhat oblong in shape and in some species carry a slight amount of fine pubescence. The mesocoxal cavities are formed by the mesosternum on the inside and anterior border, and by the mes-episternum on the outside. Metathorax fully twice as long as the mesothorax, sternum large and very convex in the greater part of the species, slightly prolonged behind between the hind coxae. The sternum is rarely very pubescent although in some species of the subfasciata group such is the case. The metapleurae are divided as the mesopleurae, the episternum equal in length to the sternum but tapering posteriorly to a sharp point. Epimeron of equal length but widening posteriorly so that the inner posterior angle touches the outer posterior angle of the sternum. Taken together as the pleuron, the width is slightly less than the length and the general shape is rectangular. In many of the species, the epimeron is densely pubescent with rather short





silvery, sometimes scale-like hairs. Metacoxal cavities very transverse, bounded by the metasternum and just reaching the posterior margin of the epimeron. Wings quite large, membranous, when not in use folded beneath the elytra. Elytra of various shapes and sizes. In the species of the unifasciata group, they completely cover the abdomen and are bent downward at the tips. The suture is closed the entire length. In the case of subfasciata and its allies, the elytra nearly reach the tip of the abdomen but do not turn down nor is the suture closed the entire length. A third type is that of pusilla where the elytra are widely dehiscent at the suture and are very short, hardly reaching the first abdominal segment. The lateral margins and apices of most of the species bear a more or less distinct serration. In two species, aegra Newm. and brunnea (an unpublished new species from Kansas) the elytra are long but absolutely truncated at the tips. This does not obtain in any other species of the genus. Abdomen of six subequal segments, more or less visible from above. The first ventral segment is slightly longer than the following segment, second through sixth equal. Pubescence sparse but evenly distributed. The terminal ventral segment is indicative of the sex of the individual. In the males, this segment is divided into two cornu or horns which serve as claspers during the process of copulation. When the segment is completely divided, the preceeding ventral is rather deeply emarginate. In other cases the final is divided for but half its length. In the female, the terminal ventral is entire and forms the lower half of a tube leading to the sexual organs. The terminal dorsal segment is hinged so that it can be drawn down to completely close the external opening.





The legs are long and thin, typically formed for an insect which is very active. Anterior coxae large, conical and prominent, middle coxae smaller and more globose while the posterior are very narrow and transverse. Other parts of the legs are similar for the three pairs. Trochanters small, oval. Femora long, fusiform, attached to the sides of the trochanters, the posterior femora often reaching the tip of the abdomen. Tibiae thin, slightly shorter than the femora. Tarsi composed of five segments. The first segment is small and is covered from above by the second which is somewhat larger. Third and fourth similar to the second and like the second carry a membranous appendage on the under side. Last segment thin, not appendaged, usually twice as long as the fourth and bearing at its tip the tarsal claws. Claws divaricate, each carrying at the base a stout, broad appendage. Tips thin and sharp. In the species with the prothorax decidedly longer than broad, the claws do not have the basal tooth, but are slender for their entire length.

From the other genera of the subfamily, this genus may be known by the following characters, incorporated in this key:

- Antennae without terminal club . . . . . No New England genera.
- Antennae with terminal club . . . . . 1.
- 1. Club formed of three segments . . . . . No New England genera.
- Club formed of two segments . . . . . 2.
- 2. Species inhabiting the old world . . . . . Not treated.
- Species inhabiting the new world . . . . . Hydnocera Newm.

Hydnocera Newman.

Hydnocera. Newman, Entom. Mag., V, 379 (1838); Spinola, Rev. Mag. Zool., 75 (1841); Klug, Clerii, 311 (1842); Spinola, Mon. Clérites, II, 39 (1844); Leconte, Ann. Lyc. Nat. Hist. New York, V, 26, (1852); Lacordaire, Gen. Col. IV, 471 (1857);





Leconte, Class. Col. N. Amer., 3, 196 (1862); Gorham, Trans. Ent. Soc. Lond., 260, (1877); Gorham, Biol. Centr. Amer., Col. III, 168 (1883).

Clerus. Say, Journ. Acad. Phila. III, 192 (1823); Germar, Ins. Spec. Nov., 80 (1824).

Trichodes. Say, Boston Journ. I, 164 (1835).

Theano. Castelnau, Ann. Soc. Ent. Fr., (2), I, 33 (1843).

The insects of this genus are found mainly on the foliage of the various species of Cupuliferae, especially on that of the genera *Carya* and *Quercus*. Some may be beaten from certain species of *Salix* and some from the herbage and shrubs along roadsides and watercourses. Although nothing of importance has been published concerning their life-histories, all are probably predaceous or parasitic. The latter inference is supported by the fact that certain species have been reared from the galls of some hymenopterous insect. In distribution the species are known to occur from Manitoba, Canada, throughout the United States, Mexico and Central America and as far south as Buenos Ayres, Argentina. Certain of the species are prone to variation in color but this variation is not coordinate with the geographical range of the species. Other species are constant in color and form in all cases. *H. lecontei* Wolc. as it is now accepted, varies considerably in size, pubescence and even form, and it is quite possible that more than one species are included under this head. The genus seems to be comparatively young in history, as the structure of the internal genitalia is practically identical in species widely differing in their external appearance. There is one fossil example, *H. wolcotti* Wickham, described from the well known Mio-





cene deposits at Florrissant, Colorado.

Of the enemies and economic importance of these interesting insects little is known. No internal parasites have been reported as reared from any of the species but there are certainly birds and other animals which do prey upon them. The dissection of the stomach and intestine of a specimen of *Hyla versicolor* Lec. (Batrachia: Salientia: Hylidae), collected at Wilbraham, Mass. while sweeping the young hickory, showed an elytron of the species *pallipennis* Say, a form very abundant in that region. So far as is known, the early stages of these beetles are passed in the twigs of hickory or other tree, where the larvae feed on the young woodborers and Scolytids. In this way, the species are of undoubted value to the forester.

The eleven New England species may be readily separated from one another by the following table :

Prothorax distinctly longer than broad; tarsal claws without basal tooth, . . . . .	1.
Prothorax not distinctly longer than wide; tarsal claws with a broad basal tooth, . . . . .	2.
1. Upper parts entirely black; legs rufotestaceous, sometimes infusate at the knees, . . . . .	<i>tabida</i> Lec. (p. )
Upper parts brownish black; elytra usually with pale markings; legs pale yellowish, never darker, . . . . .	
. . . . .	<i>curtipennis</i> Newm. (p. )
2. Prothorax equilateral, or slightly longer than broad, . . . . .	3.
Prothorax broader than long, . . . . .	6.
3. Elytra with side margins parallel to apical fourth, . . . . .	
. . . . .	<i>commixta</i> n. sp. (p. )
Elytral margins convergent posteriorly, . . . . .	4.
4. Head and thorax with pale markings, . . . . .	<i>verticalis</i> Say. (p. )
Head and thorax aeneous black, . . . . .	5.
5. Elytra with pale markings, . . . . .	<i>pallipennis</i> Say. (p. )
Elytra without definite pale spots, . . . . .	<i>subaenea</i> Spin. (p. )





6. Elytra slightly abbreviated, tips serrate, lecontei Wolc. (p. )  
Elytra entirely covering the abdomen, . . . . . 7.
7. Sides of elytra parallel, . . . . . 8.  
Sides of elytra convergent toward apices, . . . . . 9.
8. Elytra with a transverse median bar of white pubescence, . . .  
. . . . . unifasciata Say. (p. )  
Elytra without the transverse bar; humeri sometimes red-  
dish, . . . . . humeralis Say. (p. )
9. Head finely, densely punctate, subfasciata fraterna Wolc. (p. )  
Head coarsely, less densely punctate, . . maritima Wolc. (p. )

\*\*\*\*\*

1. *Hydnocera tabida* Leconte. (center).

1849. *Hydnocera tabida* Leconte, Ann. Lyc. Nat. Hist. N. Y.,  
V, 29.

1853. *Hydnocera tabida* Melsheimer, Catalogue, 83.

1866. " *tabida* Leconte, List Col. N. A.,

1895. " *tabida* Wickham, Can. Ent., XXVII, 252.

1910. " *tabida* Wolcott, In Blatchley's Col. Ind. p. 857.

Location of type : Museum Comparative Zoology, Harvard University.

~~*H. tabida* Lec.~~ -- Very elongate; brownish black; antennae, mouth-  
parts pale, major parts of legs rufotestaceous. Head black, sur-  
face densely alutaceous and sparsely punctured, pubescence fine,  
depressed and sparse. Eyes large, convex and prominent. Thorax  
longer than broad, (measured ratio of 36 - 30), nearly cylindri-  
cal, lateral dilation weak, apical impression nearly obsolete,  
surface alutaceous with a few scattered punctures bearing fine  
hairs. Scutellum black, polished. Elytra long and narrow, coarsely  
and closely but evenly punctured, entirely black, pubescence long,  
sparse and erect. Apices separately rounded, obsoletely serrate,  
dehiscent at suture. Under parts black, alutaceous, sparsely pu-  
bescent. Legs rufotestaceous, middle and hind tibiae, and hind  
femora at apices are usually darker. Length : 4 - 5 mm.





An elongate species, distinguished from the darker forms of *curtipennis* Newm. by the darker legs and the proportionately longer elytra. The species is widely distributed, reported from Canada, along the Atlantic coast and as far west as Kansas and Nebraska. There is but one New England record, Mt. Washington, White Mountains, New Hampshire (Slosson). Usually taken in sweepings from meadow land.

*Hydnocera curtipennis* Newman.

- 1840. *Hydnocera curtipennis* Newman, Ann. Mag. Nat. Hist. IV.,  
n.s., 364.
- 1842. *Hydnocera curtipennis* Klug, Clerii, 390.
- 1844. " *longicollis* Ziegler, Proc. Acad. Nat. Sci.  
Phila., II, 44.
- 1849. *Hydnocera curtipennis* Leconte, Ann. Lyc. Nat. Hist. N. Y.,  
V, 29.
- 1849. *Hydnocera longicollis* Leconte, loc. cit., 30.
- 1853. " *curtipennis* Melsheimer, Catalogue, 83.
- 1853. " *longicollis* Melsheimer, loc. cit., 83.
- 1866. " *curtipennis* = *verticalis* Leconte, List Col. N.  
A.
- 1866. *Hydnocera longicollis* Leconte, List Col. N. A.
- 1895. " *longicollis* Wickham, Can. Ent., XXVII, 252.
- 1903. " *curtipennis* = *verticalis* Schenkling, Gen. Ins.,  
Fasc. 13, 93.
- 1903. *Hydnocera longicollis* Schenkling, loc. cit., 93.
- 1906. " *curtipennis* = *verticalis* Schenkling, Deutsch.  
Ent. Zeit., 303.
- 1910. *Hydnocera longicollis* Wolcott, In. Statchley's Col. Ind.





Location of type : unknown to the author.

~~*H. curtipennis* Newm.~~ - Very elongate; aeneous black; antennae, mouth-parts, legs and sometimes part of elytra pale. Head black, shining, sparsely and finely punctate, pubescence sparse. Eyes large, convex, slightly prominent. Thorax distinctly longer than wide, (34 - 26), lateral dilation weak, apical and basal impressions distinct, surface alutaceus with a few scattered hairs, each hair arising in a puncture. Elytra abbreviated, attenuate, humeri distinct, punctures coarse, not confluent, margins serrate toward tips. Color brownish black, usually with a pale maculation extending from the middle of the base, along the median line on each elytron for a greater or less distance. Pubescence uniform, sparse, erect. Under parts aeneous black, pleuras rugose, median parts smooth, minutely and sparsely punctured. Legs pale with long hairs. Length : 3.5 - 5 mm.

A widely distributed species, represented from New England by the following localities : Mass., West Springfield, Springfield, Wilbraham., also from New Hampshire, Hampton, (June 28). Separated from *verticalis*, which it resembles superficially, by the toothed claws and the very much longer thorax. From *tabida*, it is separated by the characters given under that species.

*Hydnocera pallipennis* Say.

1825. *Clerus pallipennis* Say, Journ. Acad. Nat. Sci. Phila., V, 176.

1837. *Hydnocera serrata* Newman, Ent. Mag. Lond., V, 380.

1837. *Phyllobaenus 4-maculatus* Dejean, Catalogue, 3rd. ed., 127.





1837. *Phyllobaenus limbatus* ♂ Dejean, loc. cit., 127.  
 1837.       "       *transversalis* ♀ Dejean, loc. cit., 127.  
 1837.       "       *aequinoctialis* (var), Dejean, loc. cit., 127.  
 1840. *Hydnocera serrata* Newman, Ann. Mag. Nat. Hist.; IV, n.s.,  
       363.  
 1842. *Hydnocera serrata* Klug, Clerii, 390.  
 1842. *Clerus pallipennis* Klug, loc. cit., 390.  
 1844. *Hydnocera serrata* Spinola, Mon. Clér., II, 47.  
 1844.       "       *limbata* Spinola, loc. cit., 49.  
 1844.       "       *suturalis* Spinola, loc. cit., Planches, 39, 6.  
 1849.       "       *pallipennis* Leconte, Ann. Lyc. Nat. Hist. N. Y.,  
       V, 28.  
 1853. *Hydnocera pallipennis* Melsheimer, Catalogue, 83.  
 1866.       "       *pallipennis* Leconte, List Col. N. A.,  
 1866.       "       *suturalis* Leconte, loc. cit.,  
 1895.       "       *pallipennis* Wickham, Can. Ent., XXVII, 251.  
 1910.       "       *pallipennis* Wolcott, In Blatchley's Col. Ind.,  
       356.

Location of type : lost with the Say collection.

~~*H. pallipennis* Say.~~   Elongate; aeneous black; elytra pale with  
 margins, suture, apices, median transverse band and humeri dark,  
 antennae, mouthparts and legs testaceous. Head black, bronzed,  
 closely and finely punctured, vertex wrinkled, pubescence sparse.  
 Eyes large convex and prominent. Thorax broader than long, (32 -  
 35), coarsely and confluentlly punctured except for disk which is  
 finely alutaceous. Anterior and basal impressions strong, sides  
 dilated strongly just before the middle, lateral foveae double,





shallow but very evident, pubescence sparse. Scutellum black with silvery pubescence. Elytra slightly shorter than the abdomen, dehiscent at the suture, punctures coarse and a little confused toward tips, which are rounded separately and strongly serrate and somewhat tumid. Color testaceous, margins, humeri, suture, apices and a median transverse band dark brownish black, these markings increasing in size at times to nearly obscure the pale color. Under parts aeneous black, mesopleurae strongly reticulate, metapleurae finely punctured and covered with silvery, silken pubescence. Legs testaceous, apices of femora and tibiae of middle and posterior legs dark. Length : 3.5 - 4.8 mm.

A very abundant species in the eastern part of the United States, but reported from nearly all the states of the Union. The New England records are : New Hampshire, Mt. Washington, Massachusetts, Tyngsboro, Mt. Tom, Springfield, West Springfield, Westfield, Agawam. Also from Maine (no definite locality cited).

Hydnocera verticalis Say.

1835. *Trichodes verticalis* Say, Bost. Journ. Nat. Hist., I, 164.

1842. *Clerus brachypterus* Klug, *Clerii*, 313.

1842. " *verticalis* Klug, loc. cit., 369.

1844. *Hydnocera lineaticollis* Spinola, Mon. Clér., II, 51.

1849. " *verticalis* Leconte, Ann. Lyc. Nat. Hist. N. Y., V, 29.

1853. *Hydnocera verticalis* Melsheimer, Catalogue, 83.

1866. " *verticalis* Leconte, List Col. N. A.,





1906. *Hydnocera verticalis* Schenkling, Deutsch. Ent. Zeit.,  
303.

1910. *Hydnocera verticalis* Wolcott, In Blatchley's Col.  
Ind., 857.

Location of type : lost in the Say collection.

~~*Hydnocera verticalis* Say.~~ - Elongate; aeneous black, front of head, antennae, mouth-parts, part of legs and usually two vittae on thorax pale. Head shining, sparsely and minutely punctured, vertex dark, sides pale. Pubescence not dense, pale, suberect. Thorax slightly broader than long, (25 - 27), black, disk smooth, sides finely rugose. Apical and basal impressions weak though distinct, lateral dilation weak, pubescence long, sparse, erect. Often with two pale vittae extending from the pale sides of the head nearly to the base. Elytra abbreviated, dehiscent at suture, usually colored as in *curtipennis* Newm., punctures coarse, evenly distributed, apices serrate, slightly tumid. Under parts black, side pieces more coarsely punctured than median, pubescent. Legs in the darker specimens more or less infuscate on femora and tibiae, in the lighter specimens nearly entirely pale. Length : 3 - 4 mm. ♂, ♀.

This species is sometimes confused in collections with *curtipennis* Newm., but is distinguished from that species by the tooth on the tarsal claw and by the nearly equilateral thorax. It is widely distributed and has been reported from nearly all sections of the country. From New England, it is known as follows: New Hampshire, White Mts.; Vermont, St. Albans; Massachusetts, Tyngsboro, Springfield, Wilbraham, Chicopee, Mt. Tom.





Hydnocera subaenea Spinola.

1844. *Hydnocera subaenea* Spinola, Mon. Clér., II, 51.  
1844. " *steniformis* Spinola, loc. cit., Planches, 40, 2.  
1853. " *subaenea* Melsheimer, Catalogue, 82.  
1866. " *subaenea* Leconte, List Col. N. A.,

Location of type : unknown to the author.

~~*Hydnocera subaenea Spin.*~~—Elongate, parallel; aeneous; elytra and legs piceous, antennae, mouth-parts, knees, front tibiae and all tarsi paler. Head finely and sparsely punctured, finely rugose on vertex, pubescence sparse. Eyes large and prominent. Thorax broader than long, (28 - 35), sparsely punctured, finely rugose on sides, disk smooth and shining. Lateral dilation weak, foveae present though indistinct, apical impression fine, deep and distinct, basal a rounded groove. Pubescence sparse, erect and dark. Scutellum black, sparsely pubescent. Elytra nearly as long as abdomen, parallel, slightly debiscent at suture, tips bluntly rounded and coarsely serrate, surface densely punctured, the punctures shallow and indistinct, becoming confused toward apices, tips very slightly tumid. Color piceous, slightly paler across base. Under parts aeneous black, sparsely and very finely punctured, sparsely pubescent. Legs piceous, knees, tarsi and anterior tibiae paler. Length : 3.8 mm. ♂, ♀

This appears to be a very rare species, described by Spinola from "L'Amerique septentrionale (Dr. Le Conte)." There are two specimens in the Wolcott collection which fit the original description very closely and are undoubtedly correctly placed. These specimens are from "Mass" and "Tyngs(boro) Mass" respectively. Not recorded from any other locality. (cf. *H. lecontei* Wolc.).





Hydnocera commixta n. sp.

Location of type : In the collection of Mr. A. B. Wolcott, Chicago, Ill., paratype (with two other specimens) in the Blanchard collection, Harvard University.

~~Hydnocera commixta n. sp.~~ - Moderately elongate; aeneous, elytra black, legs piceous, anterior and middle tibiae, all tarsi, antennae and mouth-parts paler. Head aeneous, finely and densely punctured, pubescence sparse and erect, eyes large and convex. Thorax broader than long, (37 - 42), lateral dilations moderate, foveae shallow but conspicuous, apical impression weak, broadly V-shaped, apex of V directed posteriorly, basal impression distinct; surface coarsely and not sparsely punctured, disk nearly smooth. Scutellum black, sparsely pubescent. Elytra nearly as long as abdomen, slightly convergent behind, tips obliquely truncate, coarsely serrate except on truncation, tumid. Surface coarsely confluent punctured, pubescence sparse, equally distributed, suberect, pale. Color uniform black. Under parts slightly aeneous black, mesosternum coarsely punctate, metasternum finely rugose, metapleurae finely punctate. Sparsely pubescent. Legs piceous, anterior and middle tibiae and all tarsi paler. Length : 4.5 mm. ♂.

Type labelled "N (ew) Y(ork), Sherman." Other specimens "Mass" and "Drac(ut, Mass) - 7/28-'10." Quite distinct from any other New England species but somewhat of the appearance of a small specimen of H. tristis Schaeffer.

Hydnocera maritima Wolcott.

1910. *Hydnocera maritima* Wolcott, Ent. News, XXI, 321.





Location of type : in the Wolcott collection, Chicago, Ill.

~~Hydnocera caritina Wolcott~~.-- Elongate; olive green; antennae, mouth-parts, tibiae and tarsi paler. Head aeneous, more coarsely and less densely punctured than in subfasciata fraterna Wolc., sparsely pubescent. Eyes moderately prominent. Thorax broader than long, (43 - 50), aeneous, coarsely and densely punctate, apical impression deep, distinct, basal wider but fully as deep, disk with a short longitudinal sulcus just posterior to anterior impression. Lateral dilation strong, foveae deep and distinct. Scutellum with dense white pubescence on median line. Elytra elongate, slightly dehiscent at suture, apices separately rounded and irregular in outline. Pubescence white, sparse but fairly evenly distributed. Color uniform olivaceous. Under parts greenish, black, sparsely and finely punctured, finely rugose at sides. Posterior half of metapleurae with dense whitish pubescence. Legs greenish, tibiae and tarsi paler, the hind tibiae slightly darker. Length : 5 - 5.5 mm. ♂

Very close to some forms of lecontei Wolc. but distinct in the form of the elytra. In some specimens the longitudinal sulcus of the thorax is missing. The type locality is Truro, (Cape Cod), Mass., the type collected on beach grass by Mr. Frederick Blanchard in June, 1892. Specimens have been seen from Long Island. The species is probably restricted to the coastal regions in distribution.

Hydnocera lecontei Wolcott.

1849. Hydnocera subaenea Leconte, Ann. Lyc. Nat. Hist. N. Y.,  
V, 26.





1912. *Hydnocera lecontei* Wolcott, Bull. Univ. Iowa, VI, 62.

Location of type : in the Leconte collection at Harvard University.

~~*Hydnocera lecontei* Wolcott.~~ - Moderately robust, form of subfasciata Lec.; metallic black to greenish; antennae, mouth-parts and tarsi piceous. Head finely punctured, quite densely, eyes convex and prominent. Thorax broader than long, (41 - 52), apical and basal impressions distinct; lateral foveae deep, distinct; lateral dilations moderately strong. Surface closely and moderately finely punctured, the punctures becoming scarce on the disk. Scutellum black, pubescent. Elytra slightly shorter than body, suture nearly closed, tips separately rounded and sparsely set with fine teeth. Pubescence moderately dense, often forming a rather indistinct post-median cross fascia. Surface densely punctate, the punctures becoming very confused at about apical third. Color uniform greenish black, sparsely or at times decidedly metallic. Underparts greenish black, sparsely punctured and pubescent. Legs greenish, tarsi and occasionally front tibiae piceous. Length : 3.8 - 5 mm. ♂, ♀.

This is the species referred to as subaenea Spin. by Dr. Leconte in his Synopsis. It has a very wide distribution and the extreme forms differ greatly. The New England specimens are fairly typical and are from the following localities : Maine, Wales, Mammoth and Paris; New Hampshire, Mt. Washington; Vermont, Brattleboro; Massachusetts, Tyngsboro, Nantucket Island, Mt. Tom, Wilbraham, Chicopee. Easily separated from subfasciata fraterna Wolc., its nearest ally, by the dark tibiae.





Hydnocera humeralis Say.

1823. *Clerus humeralis* Say, Journ. Acad. Nat. Sci. Phila.,  
III, 192.
1824. *Clerus humeralis* Germar, Ins. Nov., I, 80.
1840. " *humeralis* Klug, Abh. Berl. Akad., 312.
1840. *Hydnocera humeralis* Newman, Ent. Mag. Lond., IV, 362.
1844. " *humeralis* Spinola, Mon. Clér., II, 44.
1849. " *humeralis* Leconte, Ann. Lyc. Nat. Hist. N. Y.,  
V, 27.
1849. *Hydnocera difficilis* Leconte, loc. cit., 27.
1849. " *cyanescens* Leconte, loc. cit., 28.
1853. " *humeralis* Melsheimer, Catalogue, 83.
1853. " *cyanescens* Melsheimer, loc. cit., 83.
1866. " *humeralis* Leconte, List N. A. Col.,
1866. " *difficilis* Leconte, loc. cit.,
1895. " *humeralis* Wickham, Can. Ent., XXVII, 251.
1895. " *difficilis* Wickham, loc. cit., 251.
1906. " *difficilis* = var. of *humeralis* Schenkling,  
Deutsch. Ent. Zeit., 303.
1910. *Hydnocera humeralis* Wolcott, In Blatchley's Col. Ind., 856.

Location of type : lost in the Say collection.

~~*Hydnocera humeralis* Say.~~ Oblong, parallel; bluish black, humeri varying from reddish to black; antennae, mouth-parts, and part of legs piceous. Head black, shining, punctures deep and distinct but sparse, vertex transversely rugose; sparsely pubescent. Eyes convex, prominent. Thorax broader than long, (39 - 47), sides broadly dilated, apical impression shallow, on disk becoming several lines, basal shallow, broad and single. Surface





coarsely and irregularly rugose, the grooves more or less connecting the punctures, disk nearly smooth, sides very rough. Lateral foveae double, large and shallow. Scutellum black. Elytra entirely covering the abdomen, coarsely and evenly punctured, pubescence uniform and evenly distributed. Suture closed, tips of elytra together rounded and coarsely serrate. Color varying from blue black with the humeri and sometimes apical third rufous to entirely<sup>1</sup> blue black. Occasionally a specimen is seen that is nearly blue. Under parts black, mesopleurae strongly reticulated, metapleurae and sternum nearly smooth, minutely and sparsely punctured. Pubescence sparse, more dense on pleurae. Legs varying as elytra from rufous with the posterior femora dark to entirely black. Length : 3.5 - 5 mm. ♂, ♀.

There seems no valid reason for the retention of the name *difficilis* Lec. for the entirely blue black forms, as the intermediate forms can be found in any good series and are quite as abundant in any region as the typical form. The species is widely distributed and is reported from the following localities in New England : Maine, Belfast, Capens; New Hampshire, Durham, Lee; Vermont, St. Albans; Massachusetts, Mt. Tom, Wilbraham, Nantucket Is.; Connecticut, Darien.

*Hydnocera subfasciata fraterna* Wolcott.

1912. *Hydnocera subfasciata* var. *fraterna* Wolcott, Bull. Univ. Iowa, VI, 61.

Location of type (cotypes) : four in the collection of Mr. A. B. Wolcott and two in the collection of Mr. J. A. Frost, Framingham, Mass., the collector of the species.





~~Hydnocera subfasciata Fraterna Wolcott.~~ - Somewhat robust, greenish; tibiae and tarsi rufous. Head finely and rather densely punctured, sparsely pubescent. Eyes large and convex. Thorax broader than long, (41 - 50), sides moderately dilated, lateral foveae strong, apical and basal impressions strong, surface closely but not coarsely punctured, punctures confused in front of apical impression. Pubescence long but not dense. Scutellum greenish, pubescent. Elytra as long as abdomen, suture nearly closed, apices obliquely rounded and finely serrate, punctuation coarse and dense, pubescence not dense, sometimes forming an indistinct transverse fascia, as in the typical form. Under parts greenish black, side-pieces of the metasternum densely punctate and pubescent. Legs greenish, tibiae and tarsi rufous. Length : 4.5 mm. ♂, ♀.

This subspecies of the western species subfasciata Lec. was described by Mr. Wolcott from specimens collected at Chatham, Mass., July 14, 1907. There are specimens in the M. A. C. collection from Marion, Mass. that are referable to this subspecies.

Hydnocera unifasciata Say.

1825. *Clerus unifasciatus* Say, Journ. Acad. Nat. Sci. Phila., V, 176.

1840. *Clerus unifasciatus* Klug, Abh. Berl. Akad., 390.

1849. *Hydnocera unifasciata* Leconte, Ann. Lyc. Nat. Hist. N. Y., V, 27.

1844. *Hydnocera punctata* Spinola, Men. Clér., II, 54.

1853.       "       *unifasciata* Melshelmer, Catalogue, 83.

1866.       "       *unifasciata* Leconte, List Col. N. A.,

1895.       "       *unifasciata* Wickham, Can. Ent., XXVII, 251.

1910.       "       *unifasciata* Wolcott, In Blatchley's Col. Ind, 856, fig.





Location of type : lost in the Say collection.

~~*Hydoscopus unifasciatus* Say.~~ -- Elongate, parallel. Black, slightly bluish, shining. Elytra with a median transverse band of white pubescence. Antennae and labial palpi testaceous. Head bluish black, finely rugulose with a nearly obsolete median impression, eyes large, convex and prominent. Thorax broader than long, (29 - 37), sparsely pubescent with long, pale hairs, apical and basal impressions strong, the basal more so. Sides dilated strongly before the middle with distinct lateral foveae, sides rugulose, disk nearly smooth. Elytra completely covering the abdomen, parallel, suture closed, at base equal in width to the head, humeri prominent and smooth, each elytron rounded on the outer angle and serrate behind the middle. Surface coarsely and densely punctured, pubescence except for the median band, sparse, long and erect. Median band narrow, reaching nearly across each elytron, of fine, dense, silvery pubescence. Under parts black, sides of thorax rugose, without pubescence. Legs piceous, hairy. Length : 3.5 mm. ♂, ♀.

A species not to be confused with any other occurring in this region. Only reported from Massachusetts (Framingham, Tyngsboro, Wilbraham, Westfield and Agawam), but undoubtedly occurring in the neighboring states.

\*\*\*\*\*

#### Explanation of Plate.

(Drawings so designated are from the type specimen.)

- Fig. 1. *H. pallipennis* Say. Terminal ventral abdominal segment, ♂.  
 " 2. *H. pallipennis* Say. " " " " ♀.






- Fig. 3. *H. pallipennis* Say. Tarsus of posterior leg. x
- " 4. *H. purtipennis* Newm. Tarsal claw of posterior leg. x
- " 5. *H. pallipennis* Say. Tarsal claw of posterior leg. x
- " 6.g 17. Elytra of various species showing color distribution and marginal serration.
- " 6. *H. maritima* Wole. (TYPE).
- " 7. *H. unifasciata* Say.
- " 8. *H. subaenea* Spin.
- " 9. *H. tabida* Lec.
- " 10. *H. lecontei* Wole.
- " 11. *H. subfasciata fraterna* Wole. (COTYPE).
- " 12. *H. commixta* n. sp. (TYPE).
- " 13. *H. pallipennis* Say.
- " 14. *H. pallipennis* Say. (Color variety).
- " 15. *H. curtipennis* Newm.
- " 16. *H. verticalis* Say.
- " 17. *H. humeralis* Say. (Typical form).

\*\*\*\*\*





PLATE I.



Chapin del.





# INDEX.

abdomen, 9  
aegra, 3  
aequinoctialis, 16  
anatomy, 4  
antennae, 5

bicolor, 4  
brachypterus, 17

Carya, 11  
Clerini, 2  
commixta, 12, 20  
Corynetini, 2  
coxal cavities, 8  
Cupuliferae, 11  
curtipennis, 3, 7, 12, 14

Ellipotoma, 3  
elytra, 9  
Enopliini, 2  
eyes, 5

form, 4  
front, 5

galea, 6  
gular sutures, 7

head, 5  
humeralis, 3, 13, 23  
Hydnocerini, 2  
Hyla, 12

Isolemidia, 3

labial palpi, 6  
labium, 6  
labrum, 6  
lacinea, 6  
lecontei, 11, 13, 21  
legs, 10  
ligula, 6  
limbatus, 16  
lineaticollis, 17  
longa, 4  
longicollis, 14

mandibles, 6  
maritima, 13, 20  
maxillae, 6  
maxillary palpi, 6  
mesothorax, 8  
metathorax, 8

pallipennis, 3, 12, 15  
paraglossae, 7

Phyllobaenus, 2  
prothorax, 7  
punctata, 25  
pusilla, 4

Quercus, 11

rufipes, 3

Salix, 11  
sanguineus, 3  
scabra, 5  
scolytidae, 12  
serrata, 3, 15, 16  
sexual characters, 9  
size, 4  
subaenea, 12, 19  
subfasciata, 5  
subfasciata fraterna, 13, 24

tabida, 12, 13  
Theano, 3  
thorax, 7  
Tillini, 2  
transversalis, 16

unifasciata, 3, 13, 25

versicolor, 12  
vertex, 5  
verticalis, 3, 12, 19  
vestiture, 4

wings, 9  
wolcotti, 11

Zenodosus, 3



1. 1890  
2. 1891  
3. 1892  
4. 1893  
5. 1894  
6. 1895  
7. 1896  
8. 1897  
9. 1898  
10. 1899  
11. 1900  
12. 1901  
13. 1902  
14. 1903  
15. 1904  
16. 1905  
17. 1906  
18. 1907  
19. 1908  
20. 1909  
21. 1910  
22. 1911  
23. 1912  
24. 1913  
25. 1914  
26. 1915  
27. 1916  
28. 1917  
29. 1918  
30. 1919  
31. 1920  
32. 1921  
33. 1922  
34. 1923  
35. 1924  
36. 1925  
37. 1926  
38. 1927  
39. 1928  
40. 1929  
41. 1930  
42. 1931  
43. 1932  
44. 1933  
45. 1934  
46. 1935  
47. 1936  
48. 1937  
49. 1938  
50. 1939  
51. 1940  
52. 1941  
53. 1942  
54. 1943  
55. 1944  
56. 1945  
57. 1946  
58. 1947  
59. 1948  
60. 1949  
61. 1950  
62. 1951  
63. 1952  
64. 1953  
65. 1954  
66. 1955  
67. 1956  
68. 1957  
69. 1958  
70. 1959  
71. 1960  
72. 1961  
73. 1962  
74. 1963  
75. 1964  
76. 1965  
77. 1966  
78. 1967  
79. 1968  
80. 1969  
81. 1970  
82. 1971  
83. 1972  
84. 1973  
85. 1974  
86. 1975  
87. 1976  
88. 1977  
89. 1978  
90. 1979  
91. 1980  
92. 1981  
93. 1982  
94. 1983  
95. 1984  
96. 1985  
97. 1986  
98. 1987  
99. 1988  
100. 1989  
101. 1990  
102. 1991  
103. 1992  
104. 1993  
105. 1994  
106. 1995  
107. 1996  
108. 1997  
109. 1998  
110. 1999  
111. 2000  
112. 2001  
113. 2002  
114. 2003  
115. 2004  
116. 2005  
117. 2006  
118. 2007  
119. 2008  
120. 2009  
121. 2010  
122. 2011  
123. 2012  
124. 2013  
125. 2014  
126. 2015  
127. 2016  
128. 2017  
129. 2018  
130. 2019  
131. 2020  
132. 2021  
133. 2022  
134. 2023  
135. 2024  
136. 2025  
137. 2026  
138. 2027  
139. 2028  
140. 2029  
141. 2030  
142. 2031  
143. 2032  
144. 2033  
145. 2034  
146. 2035  
147. 2036  
148. 2037  
149. 2038  
150. 2039  
151. 2040  
152. 2041  
153. 2042  
154. 2043  
155. 2044  
156. 2045  
157. 2046  
158. 2047  
159. 2048  
160. 2049  
161. 2050  
162. 2051  
163. 2052  
164. 2053  
165. 2054  
166. 2055  
167. 2056  
168. 2057  
169. 2058  
170. 2059  
171. 2060  
172. 2061  
173. 2062  
174. 2063  
175. 2064  
176. 2065  
177. 2066  
178. 2067  
179. 2068  
180. 2069  
181. 2070  
182. 2071  
183. 2072  
184. 2073  
185. 2074  
186. 2075  
187. 2076  
188. 2077  
189. 2078  
190. 2079  
191. 2080  
192. 2081  
193. 2082  
194. 2083  
195. 2084  
196. 2085  
197. 2086  
198. 2087  
199. 2088  
200. 2089  
201. 2090  
202. 2091  
203. 2092  
204. 2093  
205. 2094  
206. 2095  
207. 2096  
208. 2097  
209. 2098  
210. 2099  
211. 2100  
212. 2101  
213. 2102  
214. 2103  
215. 2104  
216. 2105  
217. 2106  
218. 2107  
219. 2108  
220. 2109  
221. 2110  
222. 2111  
223. 2112  
224. 2113  
225. 2114  
226. 2115  
227. 2116  
228. 2117  
229. 2118  
230. 2119  
231. 2120  
232. 2121  
233. 2122  
234. 2123  
235. 2124  
236. 2125  
237. 2126  
238. 2127  
239. 2128  
240. 2129  
241. 2130  
242. 2131  
243. 2132  
244. 2133  
245. 2134  
246. 2135  
247. 2136  
248. 2137  
249. 2138  
250. 2139  
251. 2140  
252. 2141  
253. 2142  
254. 2143  
255. 2144  
256. 2145  
257. 2146  
258. 2147  
259. 2148  
260. 2149  
261. 2150  
262. 2151  
263. 2152  
264. 2153  
265. 2154  
266. 2155  
267. 2156  
268. 2157  
269. 2158  
270. 2159  
271. 2160  
272. 2161  
273. 2162  
274. 2163  
275. 2164  
276. 2165  
277. 2166  
278. 2167  
279. 2168  
280. 2169  
281. 2170  
282. 2171  
283. 2172  
284. 2173  
285. 2174  
286. 2175  
287. 2176  
288. 2177  
289. 2178  
290. 2179  
291. 2180  
292. 2181  
293. 2182  
294. 2183  
295. 2184  
296. 2185  
297. 2186  
298. 2187  
299. 2188  
300. 2189  
301. 2190  
302. 2191  
303. 2192  
304. 2193  
305. 2194  
306. 2195  
307. 2196  
308. 2197  
309. 2198  
310. 2199  
311. 2200  
312. 2201  
313. 2202  
314. 2203  
315. 2204  
316. 2205  
317. 2206  
318. 2207  
319. 2208  
320. 2209  
321. 2210  
322. 2211  
323. 2212  
324. 2213  
325. 2214  
326. 2215  
327. 2216  
328. 2217  
329. 2218  
330. 2219  
331. 2220  
332. 2221  
333. 2222  
334. 2223  
335. 2224  
336. 2225  
337. 2226  
338. 2227  
339. 2228  
340. 2229  
341. 2230  
342. 2231  
343. 2232  
344. 2233  
345. 2234  
346. 2235  
347. 2236  
348. 2237  
349. 2238  
350. 2239  
351. 2240  
352. 2241  
353. 2242  
354. 2243  
355. 2244  
356. 2245  
357. 2246  
358. 2247  
359. 2248  
360. 2249  
361. 2250  
362. 2251  
363. 2252  
364. 2253  
365. 2254  
366. 2255  
367. 2256  
368. 2257  
369. 2258  
370. 2259  
371. 2260  
372. 2261  
373. 2262  
374. 2263  
375. 2264  
376. 2265  
377. 2266  
378. 2267  
379. 2268  
380. 2269  
381. 2270  
382. 2271  
383. 2272  
384. 2273  
385. 2274  
386. 2275  
387. 2276  
388. 2277  
389. 2278  
390. 2279  
391. 2280  
392. 2281  
393. 2282  
394. 2283  
395. 2284  
396. 2285  
397. 2286  
398. 2287  
399. 2288  
400. 2289  
401. 2290  
402. 2291  
403. 2292  
404. 2293  
405. 2294  
406. 2295  
407. 2296  
408. 2297  
409. 2298  
410. 2299  
411. 2300  
412. 2301  
413. 2302  
414. 2303  
415. 2304  
416. 2305  
417. 2306  
418. 2307  
419. 2308  
420. 2309  
421. 2310  
422. 2311  
423. 2312  
424. 2313  
425. 2314  
426. 2315  
427. 2316  
428. 2317  
429. 2318  
430. 2319  
431. 2320  
432. 2321  
433. 2322  
434. 2323  
435. 2324  
436. 2325  
437. 2326  
438. 2327  
439. 2328  
440. 2329  
441. 2330  
442. 2331  
443. 2332  
444. 2333  
445. 2334  
446. 2335  
447. 2336  
448. 2337  
449. 2338  
450. 2339  
451. 2340  
452. 2341  
453. 2342  
454. 2343  
455. 2344  
456. 2345  
457. 2346  
458. 2347  
459. 2348  
460. 2349  
461. 2350  
462. 2351  
463. 2352  
464. 2353  
465. 2354  
466. 2355  
467. 2356  
468. 2357  
469. 2358  
470. 2359  
471. 2360  
472. 2361  
473. 2362  
474. 2363  
475. 2364  
476. 2365  
477. 2366  
478. 2367  
479. 2368  
480. 2369  
481. 2370  
482. 2371  
483. 2372  
484. 2373  
485. 2374  
486. 2375  
487. 2376  
488. 2377  
489. 2378  
490. 2379  
491. 2380  
492. 2381  
493. 2382  
494. 2383  
495. 2384  
496. 2385  
497. 2386  
498. 2387  
499. 2388  
500. 2389  
501. 2390  
502. 2391  
503. 2392  
504. 2393  
505. 2394  
506. 2395  
507. 2396  
508. 2397  
509. 2398  
510. 2399  
511. 2400  
512. 2401  
513. 2402  
514. 2403  
515. 2404  
516. 2405  
517. 2406  
518. 2407  
519. 2408  
520. 2409  
521. 2410  
522. 2411  
523. 2412  
524. 2413  
525. 2414  
526. 2415  
527. 2416  
528. 2417  
529. 2418  
530. 2419  
531. 2420  
532. 2421  
533. 2422  
534. 2423  
535. 2424  
536. 2425  
537. 2426  
538. 2427  
539. 2428  
540. 2429  
541. 2430  
542. 2431  
543. 2432  
544. 2433  
545. 2434  
546. 2435  
547. 2436  
548. 2437  
549. 2438  
550. 2439  
551. 2440  
552. 2441  
553. 2442  
554. 2443  
555. 2444  
556. 2445  
557. 2446  
558. 2447  
559. 2448  
560. 2449  
561. 2450  
562. 2451  
563. 2452  
564. 2453  
565. 2454  
566. 2455  
567. 2456  
568. 2457  
569. 2458  
570. 2459  
571. 2460  
572. 2461  
573. 2462  
574. 2463  
575. 2464  
576. 2465  
577. 2466  
578. 2467  
579. 2468  
580. 2469  
581. 2470  
582. 2471  
583. 2472  
584. 2473  
585. 2474  
586. 2475  
587. 2476  
588. 2477  
589. 2478  
590. 2479  
591. 2480  
592. 2481  
593. 2482  
594. 2483  
595. 2484  
596. 2485  
597. 2486  
598. 2487  
599. 2488  
600. 2489  
601. 2490  
602. 2491  
603. 2492  
604. 2493  
605. 2494  
606. 2495  
607. 2496  
608. 2497  
609. 2498  
610. 2499  
611. 2500  
612. 2501  
613. 2502  
614. 2503  
615. 2504  
616. 2505  
617. 2506  
618. 2507  
619. 2508  
620. 2509  
621. 2510  
622. 2511  
623. 2512  
624. 2513  
625. 2514  
626. 2515  
627. 2516  
628. 2517  
629. 2518  
630. 2519  
631. 2520  
632. 2521  
633. 2522  
634. 2523  
635. 2524  
636. 2525  
637. 2526  
638. 2527  
639. 2528  
640. 2529  
641. 2530  
642. 2531  
643. 2532  
644. 2533  
645. 2534  
646. 2535  
647. 2536  
648. 2537  
649. 2538  
650. 2539  
651. 2540  
652. 2541  
653. 2542  
654. 2543  
655. 2544  
656. 2545  
657. 2546  
658. 2547  
659. 2548  
660. 2549  
661. 2550  
662. 2551  
663. 2552  
664. 2553  
665. 2554  
666. 2555  
667. 2556  
668. 2557  
669. 2558  
670. 2559  
671. 2560  
672. 2561  
673. 2562  
674. 2563  
675. 2564  
676. 2565  
677. 2566  
678. 2567  
679. 2568  
680. 2569  
681. 2570  
682. 2571  
683. 2572  
684. 2573  
685. 2574  
686. 2575  
687. 2576  
688. 2577  
689. 2578  
690. 2579  
691. 2580  
692. 2581  
693. 2582  
694. 2583  
695. 2584  
696. 2585  
697. 2586  
698. 2587  
699. 2588  
700. 2589  
701. 2590  
702. 2591  
703. 2592  
704. 2593  
705. 2594  
706. 2595  
707. 2596  
708. 2597  
709. 2598  
710. 2599  
711. 2600  
712. 2601  
713. 2602  
714. 2603  
715. 2604  
716. 2605  
717. 2606  
718. 2607  
719. 2608  
720. 2609  
721. 2610  
722. 2611  
723. 2612  
724. 2613  
725. 2614  
726. 2615  
727. 2616  
728. 2617  
729. 2618  
730. 2619  
731. 2620  
732. 2621  
733. 2622  
734. 2623  
735. 2624  
736. 2625  
737. 2626  
738. 2627  
739. 2628  
740. 2629  
741. 2630  
742. 2631  
743. 2632  
744. 2633  
745. 2634  
746. 2635  
747. 2636  
748. 2637  
749. 2638  
750. 2639  
751. 2640  
752. 2641  
753. 2642  
754. 2643  
755. 2644  
756. 2645  
757. 2646  
758. 2647  
759. 2648  
760. 2649  
761. 2650  
762. 2651  
763. 2652  
764. 2653  
765. 2654  
766. 2655  
767. 2656  
768. 2657  
769. 2658  
770. 2659  
771. 2660  
772. 2661  
773. 2662  
774. 2663  
775. 2664  
776. 2665  
777. 2666  
778. 2667  
779. 2668  
780. 2669  
781. 2670  
782. 2671  
783. 2672  
784. 2673  
785. 2674  
786. 2675  
787. 2676  
788. 2677  
789. 2678  
790. 2679  
791. 2680  
792. 2681  
793. 2682  
794. 2683  
795. 2684  
796. 2685  
797. 2686  
798. 2687  
799. 2688  
800. 2689  
801. 2690  
802. 2691  
803. 2692  
804. 2693  
805. 2694  
806. 2695  
807. 2696  
808. 2697  
809. 2698  
810. 2699  
811. 2700  
812. 2701  
813. 2702  
814. 2703  
815. 2704  
816. 2705  
817. 2706  
818. 2707  
819. 2708  
820. 2709  
821. 2710  
822. 2711  
823. 2712  
824. 2713  
825. 2714  
826. 2715  
827. 2716  
828. 2717  
829. 2718  
830. 2719  
831. 2720  
832. 2721  
833. 2722  
834. 2723  
835. 2724  
836. 2725  
837. 2726  
838. 2727  
839. 2728  
840. 2729  
841. 2730  
842. 2731  
843. 2732  
844. 2733  
845. 2734  
846. 2735  
847. 2736  
848. 2737  
849. 2738  
850. 2739  
851. 2740  
852. 2741  
853. 2742  
854. 2743  
855. 2744  
856. 2745  
857. 2746  
858. 2747  
859. 2748  
860. 2749  
861. 2750  
862. 2751  
863. 2752  
864. 2753  
865. 2754  
866. 2755  
867. 2756  
868. 2757  
869. 2758  
870. 2759  
871. 2760  
872. 2761  
873. 2762  
874. 2763  
875. 2764  
876. 2765  
877. 2766  
878. 2767  
879. 2768  
880. 2769  
881. 2770  
882. 2771  
883. 2772  
884. 2773  
885. 2774  
886. 2775  
887. 2776  
888. 2777  
889. 2778  
890. 2779  
891. 2780  
892. 2781  
893. 2782  
894. 2783  
895. 2784  
896. 2785  
897. 2786  
898. 2787  
899. 2788  
900. 2789  
901. 2790  
902. 2791  
903. 2792  
904. 2793  
905. 2794  
906. 2795  
907. 2796  
908. 2797  
909. 2798  
910. 2799  
911. 2800  
912. 2801  
913. 2802  
914. 2803  
915. 2804  
916. 2805  
917. 2806  
918. 2807  
919. 2808  
920. 2809  
921. 2810  
922. 2811  
923. 2812  
924. 2813  
925. 2814  
926. 2815  
927. 2816  
928. 2817  
929. 2818  
930. 2819  
931. 2820  
932. 2821  
933. 2822  
934. 2823  
935. 2824  
936. 2825  
937. 2826  
938. 2827  
939. 2828  
940. 2829  
941. 2830  
942. 2831  
943. 2832  
944. 2833  
945. 2834  
946. 2835  
947. 2836  
948. 2837  
949. 2838  
950. 2839  
951. 2840  
952. 2841  
953. 2842  
954. 2843  
955. 2844  
956. 2845  
957. 2846  
958. 2847  
959. 2848  
960. 2849  
961. 2850  
962. 2851  
963. 2852  
964. 2853  
965. 2854  
966. 2855  
967. 2856  
968. 2857  
969. 2858  
970. 2859  
971. 2860  
972. 2861  
973. 2862  
974. 2863  
975. 2864  
976. 2865  
977. 2866  
978. 2867  
979. 2868  
980. 2869  
981. 2870  
982. 2871  
983. 2872  
984. 2873  
985. 2874  
986. 2875  
987. 2876  
988. 2877  
989. 2878  
990. 2879  
991. 2880  
992. 2881  
993. 2882  
994. 2883  
995. 2884  
996. 2885  
997. 2886  
998. 2887  
999. 2888  
1000. 2889  
1001. 2890  
1002. 2891  
1003. 2892  
1004. 2893  
1005. 2894  
1006. 2895  
1007. 2896  
1008. 2897  
1009. 2898  
1010. 2899  
1011. 2900  
1012. 2901  
1013. 2902  
1014. 2903  
1015. 2904  
1016. 2905  
1017. 2906  
1018. 2907  
1019. 2908  
1020. 2909  
1021. 2910  
1022. 2911  
1023. 2912  
1024. 2913  
1025. 2914  
1026. 2915  
1027. 2916  
1028. 2917  
1029. 2918  
1030. 2919  
1031. 2920  
1032. 2921  
1033. 2922  
1034. 2923  
1035. 2924  
1036. 2925  
1037. 2926  
1038. 2927  
1039. 2928  
1040. 2929  
1041. 2930  
1042. 2931  
1043. 2932  
1044. 2933  
1045. 2934  
1046. 2935  
1047. 2936  
1048. 2937  
1049. 2938  
1050. 2939  
1051. 2940  
1052. 2941  
1053. 2942  
1054. 2943  
1055. 2944  
1056. 2945  
1057. 2946  
1058. 2947  
1059. 2948  
1060. 2949  
1061. 2950  
1062. 2951  
1063. 2952  
1064. 2953  
1065. 2954  
1066. 2955  
1067. 2956  
1068. 2957  
1069. 2958  
1070. 2959  
1071. 2960  
1072. 2961  
1073. 2962  
1074. 2963  
1075. 2964  
1076. 2965  
1077. 2966  
1078. 2967  
1079. 2968  
1080. 2969  
1081. 2970  
1082. 2971  
1083. 2972  
1084. 2973  
1085. 2974  
1086. 2975  
1087. 2976  
1088. 2977  
1089. 2978  
1090. 2979  
1091. 2980  
1092. 2981  
1093. 2982  
1094. 2983  
1095. 2984  
1096. 2985  
1097. 2986  
1098. 2987  
1099. 2988  
1100. 2989  
1101. 2990  
1102. 2991  
1103. 2992  
1104. 2993  
1105. 2994  
1106. 2995  
1107. 2996  
1108. 2997  
1109. 2998  
1110. 2999  
1111. 3000  
1112. 3001  
1113. 3002  
1114. 3003  
1115. 3004  
1116. 3005  
1117. 3006  
1118. 3007  
1119. 3008  
11















